



THE GENUS *ENTEROMORPHA* LINK (ULVALES: CHLOROPHYTA) AT VISAKHAPATNAM COAST, INDIA

SUBHABRATA GHOSH and JAI PRAKASH KESHRI

Phycology Laboratory, Centre for Advanced Studies in Botany, The University of Burdwan, Golapbag, Burdwan-713104
West Bengal, India

Email : subhabrata_07@rediffmail.com ; jpkeshri@indiatimes.com

ABSTRACT

Enteromorpha Link is the most common green seaweed of Indian coasts and estuarine waters. The knowledge about the distribution of its species in India is however, insufficient. During an algal excursion to Visakhapatnam coast, the authors encountered following three taxa of the genus viz., *E. flexuosa* (Wulfen) J. Ag. subsp. *flexuosa*, *E. flexuosa* subsp. *paradoxa* (C.Ag.) Bliding and *E. lingulata* J.G. Ag. First and third taxa constitute new record from Visakhapatnam coast.

KEY WORDS: *Enteromorpha*, new records, Visakhapatnam coast, India

INTRODUCTION

Among the various sea weeds, reported from the coastal belts and estuarine pockets of India, *Enteromorpha* is most common. According to Krishnamurthy (2000) the genus is represented by 14 species and its infra specific taxa. Its distribution and morphological details have been worked out by various workers (Joshi & Krishnamurthy, 1972; Kale, 1966; Ramanathan, 1939; Biswas, 1932; Srinivasan, 1973; Dixit, 1968; Sarma & Suryanarayana, 1967) which has been summarized in recent checklists by Sahoo, *et al.* (2001) and Oza & Zaidi (2001). In spite of the works of various researchers our knowledge about the species distribution of the genus remains inadequate, particularly in the east coast of India. With this in view the present study was undertaken.

MATERIALS AND METHODS

Specimens were encountered from the intertidal zone of Visakhapatnam coast during a low tide status. Actually it is very important to go for collection, few hours before the time of low tide as per tide tables. This would provide more time for seaweed collection and their observation in natural condition. Samples were kept in plastic bags with detailed field notes. Herbarium sheets were prepared following the standard method (Dhargalkar & Kavlekar, 2004). Camera Lucida drawings were made from preserved specimens within a month from the date of collection using Olympus GB microscope. Some digital photomicrographs

were taken by LEICA DMLB photo microscopic system. The preserved materials and permanent slide preparations are stored in Phycology Laboratory, the University of Burdwan for future study and reference purpose. Identifications were made following Setchell & Gardner (1920), Joshi & Krishnamurthy (1972); Krishnamurthy (2000).

RESULTS AND DISCUSSION


The thallus of the genus shows a hollow monostromatic tube, simple to profusely branched, sometimes showing compressed nature; rhizoidal part attached with the substratum; cells angular and arranged irregularly within a mucilaginous sheath showing a tendency to lie in vertical series; each cell uninucleate with a single parietal chloroplast usually containing a single pyrenoid, although some species have more than one.

Following taxa were studied:

1. *Enteromorpha flexuosa* (Wulfen) J. Ag. subsp. *flexuosa* (Krishnamurthy, 2000, p. 90, fig. 13: J-L, 14: A-C)
(Plate: I, Fig. 1-4; Plate: II, Figs. 9-11)

Plants 6-11 cm long, 0.4-0.8 cm wide, attached to rocky substratum with the help of basal disc, light yellowish green, branched or unbranched (Lougheed & Stevenson, 2004), tubular stalk cylindrical, blade expanding above stalk and becoming flexuous, ending in an obtuse apex; cells polygonal in surface view, except in upper flexuous part;


PLATE - I



LEGEND OF FIGURES

Fig.1: *Enteromorpha flexuosa* subsp. *flexuosa* (External morphology), Fig. 2: *Enteromorpha flexuosa* subsp. *flexuosa* (Cells with chloroplast and pyrenoids), Fig. 3: *Enteromorpha flexuosa* subsp. *flexuosa* (Tip portion), Fig. 4: *Enteromorpha flexuosa* subsp. *flexuosa* (Tip portion showing bifurcation), Fig. 5: *Enteromorpha lingulata* (Disc like basal portion with branched above portion), Fig. 6: *Enteromorpha lingulata* (Cells with chloroplast and pyrenoids), Fig. 7: *Enteromorpha flexuosa* subsp. *paradoxa* (A part of tubular portion with cell arrangement), Fig. 8: *Enteromorpha flexuosa* subsp. *paradoxa* (External morphology).

PLATE - II



LEGEND OF FIGURES

Fig. 9: *Enteromorpha flexuosa* subsp. *flexuosa* (External morphology showing basal branching pattern), Fig. 10: *Enteromorpha flexuosa* subsp. *flexuosa* (Cell arrangement & chloroplast nature), Fig. 11: *Enteromorpha flexuosa* subsp. *flexuosa* (Tip portion showing bifurcation), Fig. 12: *Enteromorpha lingulata* (basal branching pattern), Fig. 13: *Enteromorpha lingulata* (Tip portion). Fig. 14: *Enteromorpha lingulata* (Cell arrangement with chloroplast nature), Fig. 15: *Enteromorpha lingulata* (Upper branching pattern).

arranged in linear series, 16-20 µm in length and 9.25-11 µm in breadth; uninucleate; chloroplast covering half the cell with 3-4 pyrenoids.

Collection No. : V18

Date: 24.01.2006

Distribution in India: Dwarka, Gujarat & Maharashtra (Børgesen, 1935); Malvan harbour, Bombay (Dixit, 1940); Madras [(Krishnamurthy, 1954 as *E. prolifera* var. *tubulosa* (Kuetz.) Batters); Mandapam camp, Gulf of Mannar (Subbaramaiah, 1972); Saurashtra coast, Gujarat (Parekh *et al.*, 1977); Maharashtra coast (Chauhan & Mairh, 1978; Dhargalkar *et al.*, 1980); Okha coast, Gujarat (Ohno *et al.*, 1981; Mairh *et al.*, 1985; Oza *et al.*, 1985); Karnataka coast (Agadi, 1985); Kerala coast (Nair *et al.*, 1990); Tuticorin (Kumar, 1993 a,b); Lakshadweep (Silva *et al.*, 1996).

This is the first report of the taxon from Visakhapatnam coast.

2. *Enteromorpha flexuosa* (Wulfen) J. Ag. subsp. *paradoxa* (C. Ag.) Bliding (Krishnamurthy, 2000, p. 90, fig. 13: D-F)

(Plate: I, Figs. 7 & 8)

Plants attached with the substratum or free floating, filiform, tubular, branched repeatedly; branches tapering from base and finally ending in single series of cells; cells rectangular to sub rectangular in surface view, 13-19.5 µm in length and 6.5-13.5 µm in breadth; uninucleate with chloroplast partially filled having 2-4 pyrenoids.

Collection No. : V7, V10

Date: 23.10.2006

Distribution in India: Saurashtra (Kale, 1966 as *E. plumosa* Kuetz.); Visakhapatnam, Andhra Pradesh (Joshi & Krishnamurthy, 1972); Muthupet Estuary, Tamil Nadu (Balakrishnan *et al.*, 1992 as *E. plumosa* Kuetz.); Bheemuri-Patram, Visakhapatnam (Umamaheswara Rao, 1999 as *E. plumosa* Kuetz.).

3. *Enteromorpha lingulata* J. G. Ag. (Krishnamurthy, 2000, p. 91, fig. 13: G-I)

(Plate: I, Figs. 5 & 6; Plate: II, Figs. 12-15)

Plants 3.5-8 cm long; yellowish green, attached to the substratum by a disc shaped basal portion, branched, younger branches directed towards the base and older towards apical region; lateral branches filiform, cells polygonal to sub rectangular in surface view and arranged in linear series throughout the thallus, 16-32 µm long and 13-19 µm wide, uninucleate with chloroplast partially filled having the 1-3 pyrenoids.

Collection No. : V19

Date: 23.10.2006

Distribution in India: Madras (Krishnamurthy, 1954

as *E. compressa* (L.) Grev. var. *lingulata*); Gujarat (Joshi & Krishnamurthy, 1972; Krishnamurthy, 2000).

This is the first report of the taxon from Vishakhapatnam coast.

ACKNOWLEDGEMENT

Authors express their thanks to the Head of the Department of Botany, The University of Burdwan for providing necessary laboratory facilities, to G. Mustafa, and to Prof. Pranjit Sarma for their constant encouragement and to Goutam Bala, Senior Research Fellow of Ecotaxonomy laboratory for his help in various ways.

REFERENCES

- Agadi, V.V. (1985). Distribution of marine algae in the littoral zone of Karnataka coast. In: Krishnamurthy, V. & A.G. Untawale (eds.) *Marine Plants*, Madras: *Seaweed Res. Utiln.*, pp. 35-42.
- Balakrishnan, S., Ravichandran, M. & Kaliaperumal, N. (1992). Studies on the distribution and standing crop of algae of Muthupet Estuary. Tamil Nadu. *Seaweed Res. Utiln.*, pp. 63-68.
- Biswas, K. (1926). Algal flora of Chilka lake. *Memoirs of the Asiatic Society of Bengal*, 11: 165-198, pls. 24-32.
- Børgesen, F. (1935). A list of marine algae from Bombay. *Kong. Danske. Vidensk. Selsk., Biologi. Meddel.*, pp. 64, 25 figs., X Plates.
- Chauhan, V.D. & Mairh, O.P. (1978). Report on survey of economic seaweed resources of Saurashtra coast, India. *Salt Res. Industry*, 14(2): 21-41.
- Dhargalkar, V.K., Jagtap, T.G., & Untawale, A.G. (1980). Biochemical constituents of seaweeds along the Maharashtra coast. *Ind. J. Mar. Sci.*, 9: 297-299.
- Dhargalkar, V.K. & Kavlekar, D. (2004). *Seaweeds a field manual*. National Institute of Oceanography, Dona Paula, Goa. pp. 36.
- Dixit, D.L. (1940). Algae of Malwan harbor. *J. Univ. Bombay*, 8(5): 82-98.
- Dixit, D.L. (1968). Species list of Indian marine algae-II. *J. Univ. Bombay*, 36(3 & 5 sect.B): 9-24.
- Joshi, H.V. & Krishnamurthy, V. (1972). The species of *Enteromorpha* from India. *Bot. J. Linn. Soc.*, 65: 120-121.
- Kale, S.R. (1966). On some green algae from the channels and reservoirs of Saurashtra Salt Works, Porbandar. *Curr. Sci.*, 35: 180-181.
- Krishnamurthy, V. (1954). Ecology and seasonal succession of the algae flora of a salt marsh of Madras. *J. Madras Univ.*, Section B 24: 161-178.
- Krishnamurthy, V. (2000). *Algae of India and Neighboring countries I. Chlorophyta*. Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, p. 210.
- Kumar, V. (1993a). Biochemical constituents of marine algae from Tuticorin coast. *Ind. J. Mar. Sci.*, 22: 138-140.

- Kumar, V. (1993b). Lipid content of some algae from Tuticorin coast. *Seaweed Res. Utiln.*, 16: 45-48.
- Lougheed, V.L. & Stevenson, R.J. (2004). Exotic marine macro algae (*Enteromorpha flexuosa* (Wulfen) J. Agardh) reach bloom proportions in a coastal lake of Michigan. *J. Great Lake. Res.*, 30(4): 538-544.
- Mairh, O.P., Pandey, R.S., Tewari, A., Oza, R.M. & Joshi H.V. (1985). Seasonal phenomena, growth and swarmer liberation rhythms in *Enteromorpha flexuosa* (Wulfen) J. Agardh (Chlorophyceae), occurring on the coast of Okha. *Proc. of Ind. Nat. Sci. Acad., Part B. Biol. Sci.*, 51: 381-385.
- Nair, N.B., Sobha, V., Chandran, R., Rathiamma, M., Maya, S. & Suryanarayanan, H. (1990). Algal resources of Kerala coast IX. Occurrence and relative abundance of Chlorophyceae along south west coast of India. *Seaweed Res. Utiln.*, 12: 125-135.
- Ohno, M., Mairh, O.P., Chauhan, V.D., Tewari, A., Oza, R.M., Joshi, H.V., Pandey, R.S. & P.S. Rao (1981). Mass cultivation of grass alga *Enteromorpha* on the coast of Okha, India. Report of USA Marine Biological Institute (Kochi University). 3: 51-59.
- Oza, R.M. & Zaidi, S.H. (2001). *A revised checklist of Indian Marine Algae*. CSMCRI, Bhavnagar, p. 196.
- Oza, R.M., Tewari, A., Joshi, H.V., Mairh, O.P. & Taqui Khan, M.M. (1985). Further experiments on the field, cultivation of *Enteromorpha* on the coast of Okha, Gujarat. In Krishnamurthy, V. and A.G. Untawale (eds.) *Marine Plants*, Madras: *Seaweed Res. Utiln.*, pp. 153-166.
- Parekh, R.G., Maru, L.V., & Dave, M.J. (1977). Chemical composition of green seaweeds of Saurashtra coast. *Bot. Mar.*, 20: 359-362.
- Ramanathan, K.R. (1939). The Morphology, Cytology and alternation of generation in *Enteromorpha compressa* (L.) Grev. var. *lingulata* (J.Ag.) Hauck. *Annal of Bot.*, ser. 2.3.: 375-398, 74 figs.
- Sahoo, D., Nivedita & Debasish (2001). *Seaweeds of Indian coast*. A.P.H. Publishing Corporation, New Delhi. p. 283.
- Sarma, Y.S.R.K & Suryanarayana, G. (1968). On the occurrence of *E. intermedia* Bliding in India and its cytology. *Phykos*, 6: 100-101.
- Setchell, W.A. & Gardner, N.L. (1920). *The Marine Algae of the Pacific coast of North America, Part II, Chlorophyceae*, University of California Press and Berkeley. p. 384.
- Silva, P.C., Basson P.W. & Moe, R.L. (1996). *Catalogue of the Benthic marine algae of Indian Ocean*. University of California Press, Benkley, USA. p. 1259.
- Srinivasan, K.S. (1973). *Phycologia indica* (Icones of Indian Marine Algae) vol.1. Calcutta: Botanical Survey of India. Pp. xvii + 60.
- Subbaramaiah, K. (1972). Intertidal ecology of a rocky shore at Mandapam Camp in the Gulf of Mannar. In: *Proc. 15th Indian Sci. Cong., Calcutta*, 1972, Part III, Abstract, Calcutta, p. 395.
- Umamaheswara Rao, M. (1999). Additions to the algae flora of Visakhapatnam Coast. *Phykos*, 38: 113-126.